

Maternity Education Program

Sepsis Antenatal Participant Resource Kit





Clinical Skills Development Service

Maternity Education Program

The resources developed for Maternity Education Program (MEP) are designed for use in any Queensland Health facility that care for patients/women who are pregnant/birthing or postnatal.



Developed by Sue Hampton, Midwifery Educator – Clinical Skills Development Service (CSDS) MNHHS, Dr Belinda Lowe, Obstetrician & Gynaecologist – Gold Coast University Hospital

Reviewed by Debbie McConnel, Timothy Hasted, Susan Lawton, Lynne Buetow, Janene Rattray, Jodi Bunn, Rachel Hentzschel, Christine Holm, Natasha Visser

Sepsis Antenatal – Facilitator Resource Kit

Published by Clinical Skills Development Service +61 3646 6500 Email <u>CSDS-Courses@csds.qld.edu.au</u> csds.qld.edu.au/mep

© Metro North Hospital and Health Service through the Clinical Skills Development Service 2020 All rights reserved.

Disclaimer: The content presented in this publication is distributed by the Queensland Government as an information source only. The State of Queensland makes no statements, representations or warranties about the accuracy, completeness or reliability of any information contained in this publication. The State of Queensland disclaims all responsibility and all liability (including without limitation for liability in negligence) for all expenses, losses, damages and costs you might incur as a result of the information being inaccurate or incomplete in any way, and for any reason reliance was placed on such information.

Who is this resource kit for?

This resource kit provides healthcare workers with knowledge and skills on assessing and managing maternal sepsis in the antenatal period.

Target audience

Midwifery and medical staff providing maternity care

Duration

45 mins (including setup, simulation and debrief)

Group size

Suited to small groups (6 - 8)

Learning objectives

By the end of the session, the learner should be able to:

- Recognise and respond to a clinically deteriorating patient.
- Demonstrate the clinical management of a labouring woman with sepsis.
- Assess the possible cause of the maternal sepsis and management.
- Demonstrate the clinical management of a birth and the fetus in the context of sepsis.

Supporting documents

- List of further readings
- SOMANZ Flowchart for assessment and management of sepsis in pregnancy
- Sepsis flow diagram
- CTG assessment tool



Overview

Despite an overall decline in maternal mortality in Australia, the maternal mortality rate from sepsis has increased. In the period 2008–2012, sepsis accounted for 11.4% of maternal deaths in Australia. Group A beta haemolytic streptococcal (GAS) infection is the most common pathogen, resulting in 25% of maternal deaths from sepsis in Australia. Sepsis continues to be one of the major causes of maternal mortality among Aboriginal and Torres Strait Islander women¹.

Despite significant advances, understanding of the pathobiology of sepsis remains incomplete and currently no gold standard diagnostic test exists to confirm the presence of sepsis. Sepsis is broadly defined as life-threatening organ dysfunction caused by a dysregulated host response to infection.

Early detection of sepsis is essential for appropriate multidisciplinary management to ensure the best outcomes for the mother and her baby. Septic patients may progress to develop septic shock, multi-organ failure and death.

Recognising the patient with sepsis is paramount and is the first step in appropriate assessment and management.

Screening for maternal sepsis should be performed using the omqSOFA (obstetrically modified quick sepsis related organ failure assessment) which helps account for some of the changes due to maternal physiology.

Cultures and investigations are ideally done prior to antibiotic administration, waiting for investigations should **NOT** delay therapy. Treatment for suspected sepsis should begin as soon as possible - ideally within the 'Golden Hour'.

In critically ill pregnant women with sepsis, stabilising the mother is the priority. Once in an ICU environment the obstetric team should liaise with the ICU team to plan fetal monitoring and delivery time. Continuous CTG monitoring is recommended.

Attempting delivery in the setting of maternal instability increases both maternal and fetal mortality².

Obstetric Emergency is any clinical situation involving a maternity patient where immediate medical/ midwifery assistance is required.

1 SOMANZ Guideline for the Management of Sepsis in Pregnancy 2017

2 Royal College of Obstetricians and Gynaecologists. Bacterial Sepsis in Pregnancy. Green-top Guideline No. 64a. RCOG. 2012. Available from: www.rcog.org.uk/globalassets/documents/ guidelines/gtg_64a.pdf

Further Readings

SOMANZ Guidelines for the Investigation and Management of Sepsis in Pregnancy – Society of Obstetric Medicine Australia and New Zealand

The document addresses the issue of sepsis in the peri-partum period. It contains a number of recommendations to guide clinical practice and improve patient outcomes. We have identified several key outcomes that can be audited allowing individual centres to assess their performance in implementation of these guidelines.

https://www.somanz.org/downloads/2017SepsisGuidelines.pdf

Bacterial Sepsis in Pregnancy Green-top Guideline No. 64a April 2012

The scope of this guideline covers the recognition and management of serious bacterial illness in the antenatal and intrapartum periods, arising in the genital tract or elsewhere, and its management in secondary care.

https://www.rcog.org.uk/globalassets/documents/guidelines/gtg_64a.pdf

SMFM Consult Series #47: Sepsis during pregnancy and the puerperium

The purpose of this guideline is to summarize what is known about sepsis and to provide guidance for the management of sepsis in pregnancy and the postpartum period. https://www.ajog.org/article/S0002-9378(19)30246-7/pdf

The Glasgow Structured Approach to Assessment of the Glasgow Coma Scale

The Glasgow Coma Scale provides a practical method for assessment of impairment of conscious level in response to defined stimuli.

https://www.glasgowcomascale.org/

Queensland Clinical Guideline: Intrapartum fetal surveillance

https://www.health.qld.gov.au/__data/assets/pdf_file/0012/140043/g-ifs.pdf

Readings for Scenario 1 - Standard

Queensland Clinical Guideline: Preterm labour and birth


https://www.health.qld.gov.au/__data/assets/pdf_file/0019/140149/g-ptl.pdf

Preterm prelabour rupture of membranes (PPROM)

https://www.health.qld.gov.au/__data/assets/pdf_file/0035/736964/g-pprom.pdf




Emergency Management



<complex-block>

SOMANZ Flowchart for assessment and management of sepsis in pregnancy



Sepsis (Antenatal)



Specific Management

Glasgow Coma scale video

https://www.glasgowcomascale.org/#video





Scan me on your phone

Fetal blood sampling results

| Interpretation | pH (units) | Lactate (mmol/L) |
|-------------------------------------|---------------|------------------|
| Normal | ≥ 7.25 | <u>≥</u> 4.2 |
| Borderline: Repeat in 30 minutes | ≥ 7.21 – 7.24 | 4.2 – 4.8 |
| Abnormal: Birth expediated | ≤ 7.20 | >4.8 |

| CTG Classification | | | | | | | |
|--------------------|--|--------|-----------------------------|---|------------------------------------|-----------------------------|--|
| | Classification | | Baseline | Variability | Declaration | Acceleration | Action/ Escalation |
| Normal | Low Probability Fetal compromise | GREEN | 110 -160 bpm | 6 -25 bpm | Nil | 15 bpm for 15 seconds | Nil |
| mal | Unlikely Fetal compromise | BLUE | 100 – 109 bpm | | Early or Late | Absent | Continue CTG MO/TL review |
| | <mark>Maybe</mark> Fetal compromise | YELLOW | >160 bpm or Rising | 3 – 5 bpm for >30 minutes | Complicated variable or Late | | Correct reversible causes MO/TL review |
| Abnormal | | | ≥ 2 <mark>YE</mark> L | LOW features = RED |) | | Persistent YELLOW = RED |
| | <mark>Likely</mark> Fetal compromise | RED | < 100 for > 5 minutes | < 3 bpm for > 30 minutes or Sinusoidal | | | FBS or Expedite Birth Urgent MO review |

Acronyms and abbreviations

| Term | Definition | | |
|---------|--|--|--|
| AN | Antenatal | | |
| bpm | Beats per minute | | |
| C/S | Caesarean section | | |
| CAT 1 | Category 1 | | |
| CSDS | Clinical Skills Development Service | | |
| CTG | Cardiotocograph | | |
| ECG | Electrocardiograph | | |
| FBC | Full blood count | | |
| FBS | Fetal blood sample | | |
| GAS | Group A beta haemolytic streptococcal | | |
| GCS | Glasgow coma scale | | |
| GP | General Practitioner | | |
| Hb | Haemoglobin | | |
| ICU | Intensive care unit | | |
| ieMR | Integrated electronic medical records | | |
| IVC | Intra venous cannula | | |
| Мес | Meconium | | |
| МО | Medical Officer | | |
| NAD | Nothing abnormal detected | | |
| Obs | Observations | | |
| omqSOFA | Obstetrically modified quick sepsis related organ failure assessment | | |
| ОТ | Operating theatre | | |
| PHR | Pregnancy Health Record | | |

Maternity Education Program

Sepsis Antenatal

| Presenting part | | |
|--|--|--|
| Per vagina | | |
| Queensland Maternity Early Warning Tool | | |
| Royal College of Obstetricians & Gynaecologists | | |
| Right occipital lateral | | |
| Society for Maternal – Fetal Medicine | | |
| Society of Obstetric Medicine of Australia & New Zealand | | |
| Spontaneous vaginal delivery | | |
| Team leader | | |
| Vaginal examination | | |
| | | |

Sepsis Antenatal

References

This resource kit has been inspired by the Optimus BONUS project of the Children's Health Queensland's Simulation Training Optimising Resuscitation for Kids (STORK) service. To know more information about STORK and their Optimus project, visit their website.

- Children's Health Queensland. 2020. Queensland Paediatric Emergency Care Education | CHQ. [online] Available at: <<u>https://www.childrens.health.qld.gov.au/research/education/queensland-paediatric-</u> emergency-care-education/> [Accessed 24 July 2020].
- 2. SOMANZ Guideline for the Management of Sepsis in Pregnancy 2017
- 3. Royal College of Obstetricians and Gynaecologists. Bacterial Sepsis in Pregnancy. Greentop Guideline No. 64a. RCOG. 2012. Available from: www.rcog.org.uk/globalassets/documents/ guidelines/gtg_64a.pdf

Share your feedback



Please complete our online survey and help make Maternity Education Program better.

The survey should take no more than 5 minutes to complete. Scan the QR code with your device or visit this link

https://www.surveymonkey.com/r/Z8Q398N



Sepsis Antenatal – Facilitator Resource Kit

Developed by Sue Hampton, Midwifery Educator – Clinical Skills Development Service (CSDS) MNHHS, Dr Belinda Lowe, Obstetrician & Gynaecologist – Gold Coast University Hospital

Published by Clinical Skills Development Service +61 3646 6500 Email <u>CSDS-Courses@health.qld.gov.au</u> Visit <u>csds.qld.edu.au/mep</u>